

University of Groningen

Gene-centric meta-analysis in 87,736 individuals of European ancestry identifies multiple blood-pressure-related loci

Tragante, Vinicius; Barnes, Michael R; Ganesh, Santhi K; Lanktree, Matthew B; Guo, Wei; Franceschini, Nora; Smith, Erin N; Johnson, Toby; Holmes, Michael V; Padmanabhan, Sandosh

Published in:
American Journal of Human Genetics

DOI:
[10.1016/j.ajhg.2013.12.016](https://doi.org/10.1016/j.ajhg.2013.12.016)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2014

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Tragante, V., Barnes, M. R., Ganesh, S. K., Lanktree, M. B., Guo, W., Franceschini, N., Smith, E. N., Johnson, T., Holmes, M. V., Padmanabhan, S., Karczewski, K. J., Almoguera, B., Barnard, J., Baumert, J., Chang, Y-P. C., Elbers, C. C., Farrall, M., Fischer, M. E., Gaunt, T. R., ... Keating, B. J. (2014). Gene-centric meta-analysis in 87,736 individuals of European ancestry identifies multiple blood-pressure-related loci. *American Journal of Human Genetics*, 94(3), 349-360. <https://doi.org/10.1016/j.ajhg.2013.12.016>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Gene-centric Meta-analysis in 87,736 Individuals of European Ancestry Identifies Multiple Blood-Pressure-Related Loci

Vinicius Tragante,^{1,2,98} Michael R. Barnes,^{3,98} Santhi K. Ganesh,^{4,98} Matthew B. Lanktree,⁵ Wei Guo,⁶ Nora Franceschini,⁷ Erin N. Smith,⁸ Toby Johnson,⁹ Michael V. Holmes,¹⁰ Sandosh Padmanabhan,¹¹ Konrad J. Karczewski,¹² Berta Almoguera,¹³ John Barnard,¹⁴ Jens Baumert,¹⁵ Yen-Pei Christy Chang,¹⁶ Clara C. Elbers,^{2,17} Martin Farrall,^{18,19} Mary E. Fischer,²⁰ Tom R. Gaunt,²¹ Johannes M.I.H. Gho,¹ Christian Gieger,²² Anuj Goel,^{18,19} Yan Gong,²³ Aaron Isaacs,²⁴ Marcus E. Kleber,²⁵ Irene Mateo Leach,²⁶ Caitrin W. McDonough,²³ Matthijs F.L. Meijs,¹ Olle Melander,^{27,28} Christopher P. Nelson,^{29,30} Ilja M. Nolte,³¹ Nathan Pankratz,³² Tom S. Price,³³ Jonathan Shaffer,³⁴ Sonia Shah,³⁵ Maciej Tomaszewski,²⁹ Peter J. van der Most,³¹ Erik P.A. Van Iperen,^{36,37} Judith M. Vonk,³¹ Kate Witkowska,⁹ Caroline O.L. Wong,⁹ Li Zhang,¹⁴ Amber L. Beitelshes,³⁸ Gerald S. Berenson,³⁹ Deepak L. Bhatt,⁴⁰ Morris Brown,⁴¹ Amber Burt,⁴² Rhonda M. Cooper-DeHoff,²³ John M. Connell,⁴³ Karen J. Cruickshanks,^{20,44} Sean P. Curtis,⁴⁵ George Davey-Smith,²¹ Christian Delles,⁴⁶ Ron T. Gansevoort,⁴⁷ Xiuqing Guo,⁴⁸ Shen Haiqing,³⁸ Claire E. Hastie,⁴⁶ Marten H. Hofker,^{17,49} G. Kees Hovingh,⁵⁰ Daniel S. Kim,⁵¹ Susan A. Kirkland,⁵² Barbara E. Klein,²⁰ Ronald Klein,²⁰ Yun R. Li,¹³ Steffi Maiwald,⁵⁰ Christopher Newton-Cheh,⁵³ Eoin T. O'Brien,⁵⁴ N. Charlotte Onland-Moret,^{2,17} Walter Palmas,³⁴ Afshin Parsa,³⁸ Brenda W. Penninx,⁵⁵ Mary Pettinger,⁵⁶ Ramachandran S. Vasan,⁵⁷ Jane E. Ranchalis,⁴² Paul M. Ridker,^{58,59} Lynda M. Rose,⁵⁸ Peter Sever,⁶⁰ Daichi Shimbo,³⁴ Laura Steele,¹⁰ Ronald P. Stolk,³¹ Barbara Thorand,¹⁵ Mieke D. Trip,⁶¹ Cornelia M. van Duijn,²⁴ W. Monique Verschuren,^{17,62} Cisca Wijmenga,⁶³ Sharon Wyatt,⁶⁴ J. Hunter Young,⁶⁵ Aeilko H. Zwinderman,³⁷ Connie R. Bezzina,^{66,67} Eric Boerwinkle,⁶⁸ Juan P. Casas,^{69,70} Mark J. Caulfield,⁹ Aravinda Chakravarti,⁷¹ Daniel I. Chasman,^{58,59} Karina W. Davidson,⁷² Pieter A. Doevendans,¹ Anna F. Dominiczak,¹¹ Garret A. FitzGerald,⁷³ John G. Gums,⁷⁴ Myriam Fornage,⁷⁵ Hakon Hakonarson,^{13,97} Indrani Halder,⁷⁶ Hans L. Hillege,²⁶ Thomas Illig,^{77,78} Gail P. Jarvik,⁶⁰ Julie A. Johnson,²³ John J.P. Kastelein,⁵⁰ Wolfgang Koenig,⁷⁹ Meena Kumari,⁸⁰ Winfried März,^{25,81,82} Sarah S. Murray,⁸³ Jeffery R. O'Connell,³⁸ Albertine J. Oldehinkel,⁸⁴ James S. Pankow,⁸⁵ Daniel J. Rader,⁸⁶ Susan Redline,⁸⁷ Muredach P. Reilly,⁸⁶ Eric E. Schadt,⁸⁸ Kandice Kottke-Marchant,⁸⁹ Harold Snieder,³¹ Michael Snyder,¹² Alice V. Stanton,⁹⁰ Martin D. Tobin,⁹¹ André G. Uitterlinden,⁹² Pim van der Harst,^{26,36,63} Yvonne T. van der Schouw,¹⁷ Nilesh J. Samani,^{29,30} Hugh Watkins,^{18,19} Andrew D. Johnson,⁹³ Alex P. Reiner,⁵⁶ Xiaofeng Zhu,⁶ Paul I.W. de Bakker,^{2,17,94} Daniel Levy,^{95,99} Folkert W. Asselbergs,^{1,36,96,99} Patricia B. Munroe,^{9,99,*} and Brendan J. Keating^{10,13,97,99,*}

¹Department of Cardiology, Division Heart and Lungs, University Medical Center Utrecht, 3584 CX Utrecht, the Netherlands; ²Department of Medical Genetics, Biomedical Genetics, University Medical Center Utrecht, 3584 CX Utrecht, the Netherlands; ³William Harvey Research Institute National Institute for Health Biomedical Research Unit, Barts and the London School of Medicine, Queen Mary University of London, London EC1M 6BQ, UK; ⁴Division of Cardiovascular Medicine, Departments of Internal Medicine and Human Genetics, University of Michigan Health System, Ann Arbor, MI 48109, USA; ⁵Department of Medicine, McMaster University, Hamilton, ON L8S 4L8, Canada; ⁶Department of Epidemiology and Biostatistics, Case Western Reserve University, Cleveland, OH 44106, USA; ⁷Department of Epidemiology, University of North Carolina, Chapel Hill, NC 27599, USA; ⁸Department of Pediatrics and Rady's Children's Hospital, University of California at San Diego, School of Medicine, La Jolla, CA 92093, USA; ⁹Clinical Pharmacology and Barts and The London Genome Centre, William Harvey Research Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London EC1M 6BQ, UK; ¹⁰Department of Surgery, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA; ¹¹BHF Glasgow Cardiovascular Research Centre, University of Glasgow, Glasgow G12 8TA, UK; ¹²Department of Genetics, Stanford University School of Medicine, Stanford, CA 94305, USA; ¹³Center for Applied Genomics, Abramson Research Center, The Children's Hospital of Philadelphia, Philadelphia, PA 19104, USA; ¹⁴Department of Quantitative Health Sciences, Lerner Research Institute, Cleveland Clinic, Cleveland, OH 44195, USA; ¹⁵Institute of Epidemiology II, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg 85764, Germany; ¹⁶Departments of Medicine and Epidemiology & Public Health, University of Maryland School of Medicine, Baltimore, MD 21201, USA; ¹⁷Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, 3584 CX Utrecht, the Netherlands; ¹⁸Wellcome Trust Centre for Human Genetics, University of Oxford, Oxford OX3 7BN, UK; ¹⁹Division of Cardiovascular Medicine, Radcliffe Department of Medicine, University of Oxford, Oxford OX3 9DU, UK; ²⁰Department of Ophthalmology and Visual Sciences, University of Wisconsin, Madison, WI 53726, USA; ²¹MRC Integrative Epidemiology Unit, School of Social and Community Medicine, University of Bristol, Oakfield House, Oakfield Grove, Bristol BS8 2BN, UK; ²²Institute of Genetic Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg 85764, Germany; ²³Department of Pharmacotherapy and Translational Research and Center for Pharmacogenomics, University of Florida, Gainesville, FL 32610, USA; ²⁴Genetic Epidemiology Unit, Department of Epidemiology, Erasmus Medical Center, 3015 GE Rotterdam, the Netherlands; ²⁵Medical Clinic V, Medical Faculty Mannheim, Heidelberg University, Mannheim 68167, Germany; ²⁶Department of Cardiology, University of Groningen, University Medical Center Groningen, 9700 RB Groningen, the Netherlands; ²⁷Hypertension and Cardiovascular Disease, Department of Clinical Sciences, Lund University, Malmö 20502, Sweden; ²⁸Centre of Emergency Medicine, Skåne University Hospital, Malmö 20502, Sweden; ²⁹Department of Cardiovascular Sciences, University of Leicester, Leicester LE3 9QP, UK; ³⁰NIHR Leicester Cardiovascular

Blood pressure (BP) is a heritable risk factor for cardiovascular disease. To investigate genetic associations with systolic BP (SBP), diastolic BP (DBP), mean arterial pressure (MAP), and pulse pressure (PP), we genotyped ~50,000 SNPs in up to 87,736 individuals of European ancestry and combined these in a meta-analysis. We replicated findings in an independent set of 68,368 individuals of European ancestry. Our analyses identified 11 previously undescribed associations in independent loci containing 31 genes including *PDE1A*, *HLA-DQB1*, *CDK6*, *PRKAG2*, *VCL*, *H19*, *NUCB2*, *RELA*, *HOXC@complex*, *FBN1*, and *NFAT5* at the Bonferroni-corrected array-wide significance threshold ($p < 6 \times 10^{-7}$) and confirmed 27 previously reported associations. Bioinformatic analysis of the 11 loci provided support for a putative role in hypertension of several genes, such as *CDK6* and *NUCB2*. Analysis of potential pharmacological targets in databases of small molecules showed that ten of the genes are predicted to be a target for small molecules. In summary, we identified previously unknown loci associated with BP. Our findings extend our understanding of genes involved in BP regulation, which may provide new targets for therapeutic intervention or drug response stratification.

Introduction

Blood pressure (BP) is a major, modifiable determinant of cardiovascular disease (CVD) risk.¹ Hypertension (MIM

145500) increases risk for a variety of sequelae, including coronary artery disease, heart failure (MIM 608320), stroke (MIM 601367), and peripheral vascular disease (MIM 606787).² Heritability of daytime ambulatory BP from

Biomedical Research Unit, Glenfield Hospital, Groby Road, Leicester LE3 9QP, UK; ³¹Department of Epidemiology, University of Groningen, University Medical Center Groningen, 9700 RB Groningen, the Netherlands; ³²Institute of Human Genetics, Department of Laboratory Medicine and Pathology, University of Minnesota, Minneapolis, MN 55455, USA; ³³MRC SGDP Centre, Institute of Psychiatry, London SE5 8AF, UK; ³⁴Department of Medicine, Columbia University, New York, NY 10032, USA; ³⁵UCL Genetics Institute, Department of Genetics, Evolution and Environment, University College London, Kathleen Lonsdale Building, Gower Place, London WC1E 6BT, UK; ³⁶Durrer Center for Cardiogenetic Research, ICIN-Netherlands Heart Institute, 3511 GC Utrecht, the Netherlands; ³⁷Department of Clinical Epidemiology, Biostatistics and Bioinformatics, Academic Medical Center, 1105 AZ Amsterdam, the Netherlands; ³⁸Division of Endocrinology, Diabetes, and Nutrition, University of Maryland School of Medicine, Baltimore, MD 21201, USA; ³⁹Department of Epidemiology, Tulane University, New Orleans, LA 70118, USA; ⁴⁰Brigham and Women's Hospital and Harvard Medical School, Boston, MA 02115, USA; ⁴¹Clinical Pharmacology Unit, University of Cambridge, Addenbrookes Hospital, Hills Road, Cambridge CB2 2QQ, UK; ⁴²Department of Medicine (Medical Genetics), University of Washington, Seattle, WA 98195, USA; ⁴³University of Dundee, Ninewells Hospital & Medical School, Dundee DD1 9SY, UK; ⁴⁴Department of Population Health Sciences, University of Wisconsin, Madison, WI 53726, USA; ⁴⁵Merck Research Laboratories, P.O. Box 2000, Rahway, NJ 07065, USA; ⁴⁶Institute of Cardiovascular and Medical Sciences, University of Glasgow, Glasgow G12 8TA, UK; ⁴⁷Division of Nephrology, Department of Medicine, University of Groningen, University Medical Center Groningen, 9700 RB Groningen, the Netherlands; ⁴⁸Cedars-Sinai Med Ctr-PEDS, Los Angeles, CA 90048, USA; ⁴⁹Department Pathology and Medical Biology, Medical Biology Division, Molecular Genetics, University of Groningen, University Medical Center Groningen, 9700 RB Groningen, the Netherlands; ⁵⁰Department of Vascular Medicine, Academic Medical Center, 1105 AZ Amsterdam, the Netherlands; ⁵¹Departments of Medicine (Medical Genetics) and Genome Sciences, University of Washington, Seattle, WA 98195, USA; ⁵²Department of Community Health and Epidemiology, Dalhousie University, Halifax, NS B3H 1V7, Canada; ⁵³Broad Institute of Harvard and MIT, Massachusetts General Hospital, Boston, MA 02142, USA; ⁵⁴Conway Institute of Biomolecular and Biomedical Research, University College Dublin, Belfield, Dublin 4, Ireland; ⁵⁵Department of Psychiatry/EMGO Institute, VU University Medical Centre, 1081 BT Amsterdam, the Netherlands; ⁵⁶Division of Public Health Sciences, Fred Hutchinson Cancer Research Center, Seattle, WA 98109, USA; ⁵⁷Department of Medicine, Boston University School of Medicine, Framingham, MA 02118, USA; ⁵⁸Division of Preventive Medicine, Brigham and Women's Hospital, Boston, MA 02115, USA; ⁵⁹Harvard Medical School, Boston, MA 02115, USA; ⁶⁰International Centre for Circulatory Health, Imperial College London, W2 1LA UK; ⁶¹Department of Cardiology, Academic Medical Center, 1105 AZ Amsterdam, the Netherlands; ⁶²National Institute for Public Health and the Environment (RIVM), 3720 BA Bilthoven, the Netherlands; ⁶³Department of Genetics, University of Groningen, University Medical Center Groningen, 9700 RB Groningen, the Netherlands; ⁶⁴Schools of Nursing and Medicine, University of Mississippi Medical Center, Jackson, MS 39216, USA; ⁶⁵Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA; ⁶⁶Heart Failure Research Center, Department of Clinical and Experimental Cardiology, Academic Medical Center, 1105 AZ Amsterdam, the Netherlands; ⁶⁷Molecular and Experimental Cardiology Group, Academic Medical Centre, 1105 AZ Amsterdam, the Netherlands; ⁶⁸Human Genetics Center and Institute of Molecular Medicine and Division of Epidemiology, University of Texas Health Science Center, Houston, TX 77030, USA; ⁶⁹Faculty of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, London WC1E 7HT, UK; ⁷⁰Genetic Epidemiology Group, Department of Epidemiology and Public Health, University College London, London WC1E 6BT, UK; ⁷¹Center for Complex Disease Genomics, McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA; ⁷²Departments of Medicine & Psychiatry, Columbia University, New York, NY 10032, USA; ⁷³The Institute for Translational Medicine and Therapeutics, School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA; ⁷⁴Departments of Pharmacotherapy and Translational Research and Community Health and Family Medicine, University of Florida, Gainesville, FL 32610, USA; ⁷⁵Institute of Molecular Medicine and School of Public Health Division of Epidemiology Human Genetics and Environmental Sciences, University of Texas Health Science Center at Houston, Houston, TX 77030, USA; ⁷⁶School of Medicine, University of Pittsburgh, PA 15261, USA; ⁷⁷Research Unit of Molecular Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg 85764, Germany; ⁷⁸Hannover Unified Biobank, Hannover Medical School, Hannover 30625, Germany; ⁷⁹Department of Internal Medicine II - Cardiology, University of Ulm Medical Centre, Ulm 89081, Germany; ⁸⁰Department of Epidemiology and Public Health, Division of Population Health, University College London, Torrington Place, London WC1E 7HB, UK; ⁸¹Synlab Academy, Synlab Services GmbH, Mannheim 69214, Germany; ⁸²Clinical Institute of Medical and Chemical Laboratory Diagnostics, Medical University of Graz, Graz 8036, Austria; ⁸³Department of Pathology, University of California San Diego, La Jolla, CA 92037, USA; ⁸⁴Interdisciplinary Center Psychopathology and Emotion Regulation, University of Groningen, University Medical Center Groningen, 9700 RB Groningen, the Netherlands; ⁸⁵Division of Epidemiology and Community Health, University of Minnesota, Minneapolis, MN 55454, USA; ⁸⁶Cardiovascular Institute, the Perelman School of Medicine at the University of Pennsylvania, Philadelphia, PA 19104, USA; ⁸⁷Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA 02215, USA; ⁸⁸Department of Genetics and Genomic Sciences, Mount Sinai School of Medicine, New York, NY 10029, USA; ⁸⁹Pathology and Laboratory Medicine Institute, Cleveland Clinic, Cleveland, OH 44195, USA; ⁹⁰Molecular & Cellular Therapeutics, Royal College of Surgeons in Ireland, 123 St Stephens Green, Dublin 2, Ireland; ⁹¹Department of Health Sciences, University of Leicester, Leicester LE1 7RH, UK; ⁹²Departments of Epidemiology and Internal Medicine, Erasmus Medical Center, 3015 GE Rotterdam, the Netherlands; ⁹³National Heart, Lung and Blood Institute Framingham Heart Study, Framingham, MA 01702, USA; ⁹⁴Division of Genetics, Brigham and Women's Hospital, Harvard Medical School, Boston, MA and Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA; ⁹⁵Center for Population Studies, National Heart, Lung, and Blood Institute, Framingham, MA 01702, USA; ⁹⁶Institute of Cardiovascular Science, Faculty of Population Health Sciences, University College London, London WC1E 6BT, UK; ⁹⁷Department of Pediatrics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA

⁹⁸These authors contributed equally to this work

⁹⁹These authors contributed equally to this work and are co-senior authors

*Correspondence: p.b.munroe@qmul.ac.uk (P.B.M.), bkeating@mail.med.upenn.edu (B.J.K.)

<http://dx.doi.org/10.1016/j.ajhg.2013.12.016>. ©2014 by The American Society of Human Genetics. All rights reserved.

twin studies has been estimated to be approximately 50%–60%^{3,4} and ~40% in family studies. Several studies have been performed to elucidate underlying genetic factors for BP, including genome-wide association studies (GWASs), meta-analyses, and admixture mapping.^{5–12} Efforts to dissect the genetic basis for this complex disorder have proven challenging and currently only a small portion of the total variation in BP can be explained by common genetic variants associated signals. Part of this “missing heritability” is likely to be due to as yet unknown common or low-frequency variants, and a fraction of it could be identified by increasing sample size. Increased understanding of the underlying genetics may contribute to improvement of the treatment to reduce CVD risk.

We conducted an association analysis of BP phenotypes in 87,736 individuals of European ancestry by using a gene-centric array with ~50,000 SNPs capturing variation in ~2,100 genes related to CVD, and we replicate our findings in 68,368 independent individuals. We also performed extensive bioinformatics analyses to identify regulatory regions by using the Encyclopedia of DNA Elements (ENCODE), eQTL studies, and pathway analyses. Finally, all previously reported BP-associated loci and the genes within were annotated for their suitability as drug targets.

Material and Methods

Population Characteristics

Phenotype and genotype data from 87,736 individuals of European ancestry from 36 participating studies were used for the discovery phase, and an additional 68,368 individuals of European ancestry, from 19 additional studies, were used in the replication phase. All individuals in these studies provided informed consent, and each study was approved by its own local ethics committee. Descriptions of the participating cohorts are provided in the original publications.^{5,8,11,13}

Phenotypes

Pulse pressure (PP) was defined as systolic blood pressure (SBP) minus diastolic blood pressure (DBP), and median arterial pressure (MAP) was defined as 2/3 DBP plus 1/3 SBP. Each cohort provided regression models for its respective data, adjusted for age, age-squared, body mass index (BMI), and any study-specific corrections for population substructure (based on principal components analysis). BP values were adjusted for antihypertensive drug therapy by adding a standard treatment adjustment of 15 mmHg to the SBP and 10 mmHg to the DBP values of individuals receiving treatment in the discovery and replication cohorts. These adjustments were implemented prior to the calculation of MAP and PP.

Genotyping

A total of 52,029 SNPs were included in the meta-analyses, and all SNPs were present in at least one of the three iterative versions of the Illumina HumanCVD BeadChip (also known as the “Cardio-chip” or the ITMAT-Broad-CARE [IBC] array manufactured by Illumina)¹⁴ used by all discovery cohorts. Several studies that used this array have already been published for a variety of phenotypes and disease outcomes, including coronary artery

disease,^{16,17} plasma lipids,¹⁸ BP,^{8,11,15} heart failure,¹⁹ type 2 diabetes (T2D [MIM 125853]),²⁰ and BMI (MIM 606641).²¹

Quality Control, Association, and Meta Analyses

The discovery data sets derive from three BP consortia with completely independent samples whose results have been published separately: the IBC BP consortium⁸ and the CVD-50 consortium,²² with the addition of the Beaver Dam Studies (BOSS, EHLS, BDES) consortium.¹³ The QC steps taken for SNPs and samples were similar for both consortia, namely: individuals with <90% call rate (completeness) across all SNPs were removed, as well as SNPs with <95% call rate (completeness) or SNPs causing heterozygous haploid genotype calls across all remaining individuals. SNPs were also removed if they were not in Hardy-Weinberg equilibrium ($p < 1 \times 10^{-7}$). No study included imputed data in their analyses. Further details on genotyping and QC can be found in the original publications including the BOSS, EHLS, and BDES studies.^{8,11,13}

During the meta-analysis step, SNPs with frequencies incompatible with HapMap CEU frequencies were removed (defined as >30% difference in the minor allele frequencies) for each data source separately, i.e., the meta-analysis results of IBC BP, of CVD-50, and of BOSS, EHLS, and BDES. We used inverse variance weighted meta-analysis in MANTEL²³ to obtain a fixed-effect estimate and statistical significance for each SNP. We applied genomic control²⁴ to each data set to control effects possibly resulting from population stratification or cryptic relatedness. The CARE IBC array studies, included in this meta-analysis, determined that after accounting for linkage disequilibrium (LD), the effective number of independent tests was ~20,500 for Europeans. This resulted in an experimental or “arraywide” statistical threshold of $p = 2.4 \times 10^{-6}$ to maintain a false-positive rate of 5%²⁵ and therefore we have adopted this threshold for this study. For each associated locus, the LD patterns were examined and independence between the loci identified in this study and previously published signals was verified with SNAP²⁶ ($r^2 < 0.3$).

Replication analysis was conducted in independent samples for each trait, for SNPs with association $p < 1 \times 10^{-5}$ in the discovery analysis, with a total of 68,368 individuals from the Global Blood Pressure Genetics (GBPG) consortium,¹⁰ Women’s Genome Health Study (WGHS),²⁷ and PREVENT²⁸ and LifeLines²⁹ studies. We combined discovery and replication data in a meta-analysis and accounted for testing four phenotypes (albeit highly correlated), resulting in a Bonferroni-corrected threshold of $p < 6 \times 10^{-7}$ for the combined meta-analysis of discovery and replication samples.

Definition of Associated Gene Variants and Variant Functional Analysis

The extended locus around each associated SNP was defined by identification of all SNPs showing $r^2 \geq 0.5$. Linkage disequilibrium was defined with the HaploReg tool³⁰ based on Phase I of the 1000 Genomes project. Variants showing evidence of LD with associated variants were explored for impact on gene function via Annovar³¹ and regulatory function (including eQTLs) by HaploReg³⁰ and RegulomeDB,³² which both collate data from the Encyclopedia of DNA Elements (ENCODE)³³ and nine eQTL studies.³² Associated genes were reviewed for evidence linking them to BP-related phenotypes via PubMed and “hypertension,” “cardiovascular disease,” or “vascular disease” as medical subject headings (MeSH) terms. MeSH is a controlled vocabulary created

by the National Library of Medicine (NLM) to index journal articles and books in the life sciences. All articles in MEDLINE have been annotated with MeSH by NLM curators or designees, offering a sensitive measure of correlation between biological traits and genes in the literature.³⁴ We also checked for annotation to the gene ontology term “regulation of blood pressure,” which also represents a highly curated set of genes consistently linked to blood pressure.³⁵ At a pathway level, we used GeneGo Metacore (Thomson Reuters) to construct a custom BP network based on 436 genes annotated to the “hypertension” MeSH term and gene ontology terms described above. The Metacore database is a large commercial database of curated human, rat, and mouse gene and protein interactions individually evidenced in the literature.³⁶ This allowed us to construct a core network of interacting genes that are each individually linked to blood pressure, which collectively are likely to represent a significant blood pressure gene network. We used this network as a tool to investigate direct interactions between the blood pressure gene network and the genes in the blood pressure loci reported here. By combining network data with data on gene and variant function, we were able to prioritize genes based on their level of support with respect to BP phenotypes.

Analysis of Pharmacologic Targets

We annotated genes containing variants in LD (HapMap CEU $r^2 > 0.5$) with discovered associations and analyzed information concerning potential druggability—that is, the potential for modulation of the protein target by a water-soluble small-molecule drug. Druggable proteins usually contain a defined binding pocket or active site, which could act as a site of action (pharmacophore) for an orally bioavailable small-molecule drug. We grouped proteins into four druggability classes, based on complementary annotations of the potentially druggable genome and publicly available databases of small molecules. Targets in class 1 are already drugged with a marketed drug recorded in DrugBank; class 2 have small molecules recorded in ChEMBL, which may include compounds in current development within pharmaceutical companies, and could be used as tools in animal and cellular models; class 3 are homologous to class 1 or class 2 targets; and class 4 are predicted to contain a potentially druggable pharmacophore based on de novo structure-based druggability prediction via the online available DoGSiteScorer tool,³⁷ which binds site prediction, analysis, and druggability.

Expression Quantitative Trait Loci Analysis

We identified alias identifiers for significant index SNPs by using SNAP, an online tool for LD calculations.²⁶ Further proxy SNPs displaying high LD ($r^2 > 0.8$) were identified across four HapMap builds in European ancestry samples (CEU) with SNAP. The primary SNPs and LD proxies were searched against a collated database of expression SNP (eSNP) results including the following tissues: fresh lymphocytes,³⁸ fresh leukocytes,³⁹ leukocyte samples in individuals with celiac disease (MIM 212750),²⁴ whole blood samples,^{40–43} lymphoblastoid cell lines (LCLs) derived from asthmatic children (MIM 600807),^{44,45} HapMap LCLs from three populations,⁴⁶ a separate study on HapMap CEU LCLs,⁴⁷ additional LCL population samples,^{48–51} CD19⁺ B cells,⁵² primary PHA-stimulated T cells,⁴⁸ CD4⁺ T cells,⁵³ peripheral blood monocytes,^{52,54,55} CD11⁺ dendritic cells before and after *Mycobacterium tuberculosis* infection (MIM 607948),⁵⁶ omental and subcutaneous adipose,^{40,50,57} stomach,⁵⁷ endometrial carcinomas (MIM

608089),⁵⁸ ER⁺ and ER[−] breast cancer tumor cells (MIM 114480),⁵⁹ brain cortex,^{54,60,61} prefrontal cortex,^{62,63} frontal cortex,⁶⁴ temporal cortex,^{61,64} pons,⁶⁴ cerebellum,^{61,64} three additional large studies of brain regions including prefrontal cortex, visual cortex, and cerebellum,⁶⁵ liver,^{57,66,67} osteoblasts,⁶⁸ ileum,^{57,69} lung,⁷⁰ skin,^{50,71} and primary fibroblasts.⁴⁸ MicroRNA QTLs were also queried for LCLs⁷² and gluteal and abdominal adipose.⁷³ The collected eSNP results met the criteria for association with gene expression levels as defined in the original papers. The majority of eQTLs (15/17) showed a p value of 1×10^{-6} or less. Two replication studies with more modest association were also included. We placed more value on eQTLs with low p values that showed consistent reporting across independent studies. In each case where a SNP or proxy was associated with transcript levels, we further examined the strongest eSNP for that transcript within that data set (best eSNP) and the LD between the best eSNP and BP-selected eSNPs to estimate the concordance of the BP and expression signals.

Results

Discovery Meta-analysis

In the discovery meta-analysis, four BP traits were analyzed in 87,736 individuals from 36 cohorts, as described in Table S1 available online. We analyzed SBP, DBP, MAP, and PP as continuous traits. Cohort characteristics, including age, sex, BP values, and the proportion of individuals treated with BP-lowering medications, are provided in Table S1.

Association analyses were successfully carried out for up to 48,616 SNPs that passed QC. We identified 17 SNPs that passed a suggestive discovery p value threshold of $p < 1 \times 10^{-5}$ with six SNPs showing strongest associations with SBP, i.e., lowest p value among all four traits (and one secondary association, i.e., a threshold-passing p value but with a higher p value than in another trait), two SNPs showing strongest associations with DBP (and two other secondary associations), two showing strongest associations with MAP (and two other secondary ones), and three leading associations to PP (with three secondary associations).

Replication Analyses

Replication testing was performed in 68,368 additional individuals from 19 cohorts with genome-wide SNP genotypes imputed to HapMap, only to the signals passing the threshold on discovery and previously not published. A meta-analysis of the 17 SNPs taken forward from the discovery phase with replication data showed that 11 SNPs at independent loci met our Bonferroni-corrected array-wide significance threshold of $p < 6 \times 10^{-7}$. Some of these SNPs showed association with more than one trait, which resulted in 17 previously not described associations: three loci were associated with DBP (*PDE1A* [MIM 171890], *HLA-DQB1* [MIM 604305], and *VCL* [MIM 193065]), five loci were associated with SBP (*PRKAG2* [MIM 602743], *H19* [MIM 103280], *NUCB2* [MIM 608020], *SIPA1* [MIM 602180], and *HOXC* complex [MIM 142974]), five loci




Figure 1. Overview of the Replicated Blood-Pressure-Related Findings from This Meta-analysis for Overlap with the Various Blood-Pressure-Related Traits

Abbreviations are as follows: SBP, systolic blood pressure; DBP, diastolic blood pressure; MAP, mean arterial pressure; and PP, pulse pressure.

were associated with MAP (*PDE1A*, *VCL*, *H19*, *NUCB2*, and *RELA* [MIM 164014]), and four loci were associated with PP (*CDK6* [MIM 603368], *NUCB2*, *FBN1* [MIM 134797], and *NFAT5* [MIM 604708]) (Figure 1). The association results are summarized in Table 1. We also found a suggestive association in *ERAP1* (MIM 606832) with $2.4 \times 10^{-6} > p > 6 \times 10^{-7}$, which reached array-wide significance but not Bonferroni-corrected significance (array-wide significance divided by the number of traits, four).

We compared the results of our analysis with all published associations at the time of this report (to the best of our knowledge)^{5,8–11,22,74–77} and confirmed previously reported BP associations at 27 loci with same direction of effect (at a nominal association threshold [$p < 0.05$]), out of 32 loci covered by this genotyping array, with partial sample overlap between original findings and our study (Table S2 contains all previously reported loci and our p values for these associations; each column with an X indicates a significant result was found by the referred study to the specified trait). We did not find supportive evidence of association with BP of five loci: *CACNB2* (MIM 600003), *CDH13* (MIM 601364), *C10orf107* (MIM not available), *ZNF652* (MIM 613907), and *STK39* (MIM 607648), which were found in GWAS meta-analyses. The IBC array does not contain lead SNPs or proxies for 18 of the previously reported loci identified by GWASs.

eQTLs

Several of the reported loci have significant eQTLs. rs2854275 is associated with the expression of *HLA-DRB4* in monocytes ($p = 4.82 \times 10^{-95}$) and in liver ($p = 1.31 \times 10^{-10}$) and *HLA-DRB1* (MIM 142857) in blood ($p =$

3.20×10^{-59}); rs2282978 associates with *CDK6* expression (MIM 603368) ($p = 2.6 \times 10^{-5}$); rs4746172 is in perfect LD with rs10824069, which presents a significant association with expression of the *ADK* transcript (MIM 102750) in lung ($p < 2 \times 10^{-16}$); and rs217727 is associated with increased expression of AK126915 in LCLs ($p = 2.40 \times 10^{-6}$) and skin ($p = 6.59 \times 10^{-6}$). Closer investigation of the AK126915 transcript in the UCSC Genome Browser suggests that it may represent an isoform of the mitochondrial ribosomal protein L23 (*MRPL23* [MIM 600789]) (data not shown). rs757081 is associated with expression of *NUCB2* (MIM 608020) in LCLs in asthmatics ($p = 1.56 \times 10^{-16}$) and of *SNORD14A* in LCLs ($p = 3.10 \times 10^{-5}$); rs33063 is associated with expression of another unknown transcript, LOC283970 in B cells ($p = 9.55 \times 10^{-8}$); and rs3741378 is in tight LD with several SNPs that present significant eQTL levels in lymph, liver, and other tissues. Table S4 presents the results in more detail.

Extended Locus Analysis and Variant Functional Analysis

Variants in LD ($r^2 > 0.5$) with the 11 replicated SNPs are functionally annotated in Table S5, and a nonredundant list of corresponding genes appear in Table S6. Several loci contained only one gene (*PDE1A*, *HLA-DQB1*, *CDK6*, *PRKAG2*, and *FBN1*), whereas others contained several genes (see Table S6). By comprehensive functional annotation, we reviewed the genes at each locus both for evidence of functional impact (genic and regulatory) and a rationale in BP.

Discussion

In this study we identified 11 loci associated with BP traits ($p < 6 \times 10^{-7}$) in a meta-analysis comprising 87,736 individuals of European descent. These associations were validated in a replication cohort of 68,368 individuals. The robust sample size also confirmed a nominal association ($p < 0.05$) of 27 previously reported signals, out of the 32 previously reported loci covered by this array.

Associated Loci with a Single Candidate Gene

Of the 11 loci described in this study, 5 contain only a single gene. The *PDE1A* locus contains the phosphodiesterase 1A (*PDE1A*) gene, a Ca^{2+} /calmodulin-stimulated phosphodiesterase that preferentially hydrolyzes cGMP and has an important role in regulating vascular tone and smooth muscle cell proliferation.^{78,79} The *HLA-DQB1* locus contains *HLA-DQB1* (major histocompatibility complex, class II, DQ beta 1), which encodes a class II molecule expressed in antigen-presenting cells and plays a role in the immune system by presenting peptides derived from extracellular proteins. *HLA-DQB1* alleles have been linked to essential hypertension in Chinese populations.⁸⁰ At the *PRKAG2* locus, *PRKAG2* (protein kinase, AMP-activated, gamma 2 noncatalytic subunit) is

Table 1. Significant Association Results for All Four Traits in the Meta-analysis

Locus	SNP	CHR	Pos	Effect Allele	Discovery p Values	Replication p Values	Combined Discovery + Replication		
					(n = 87,736)	(n = 68,368)	beta	SE	p Val
DBP									
PDE1A	rs16823124	2	182932372	A	1.76×10^{-6}	2.59×10^{-5}	0.262	0.041	1.95×10^{-10}
HLA-DQB1	rs2854275	6	32736406	A	2.97×10^{-8}	1.86×10^{-4}	−0.562	0.101	5.53×10^{-12}
VCL	rs4746172	10	75525848	C	2.14×10^{-5}	1.07×10^{-3}	0.230	0.043	9.14×10^{-8}
MAP									
PDE1A	rs16823124	2	182932372	A	1.10×10^{-4}	3.85×10^{-4}	0.269	0.052	1.90×10^{-7}
VCL	rs4746172	10	75525848	C	7.91×10^{-6}	8.73×10^{-3}	0.279	0.054	2.45×10^{-7}
H19	rs217727	11	1973484	A	1.63×10^{-5}	3.94×10^{-3}	0.315	0.061	2.15×10^{-7}
NUCB2	rs757081	11	17308259	G	3.23×10^{-5}	1.16×10^{-3}	0.265	0.050	1.39×10^{-7}
RELA	rs3741378	11	65165513	T	3.08×10^{-5}	2.30×10^{-3}	−0.359	0.070	2.44×10^{-7}
PP									
CDK6	rs2282978	7	92102346	C	3.82×10^{-8}	0.073	−0.268	0.049	4.51×10^{-8}
NUCB2	rs757081	11	17308259	G	4.51×10^{-7}	4.48×10^{-5}	0.321	0.050	8.94×10^{-11}
FBN1	rs1036477	15	46702218	G	6.59×10^{-6}	8.02×10^{-3}	−0.402	0.077	2.00×10^{-7}
NFAT5	rs33063	16	68197718	A	9.07×10^{-7}	0.063	0.335	0.066	4.14×10^{-7}
SBP									
PRKAG2	rs10224002	7	151045974	G	1.67×10^{-6}	0.023	0.361	0.072	5.50×10^{-7}
H19	rs217727	11	1973484	A	7.88×10^{-6}	2.95×10^{-3}	0.417	0.078	1.03×10^{-7}
NUCB2	rs757081	11	17308259	G	2.26×10^{-7}	1.28×10^{-4}	0.403	0.063	1.65×10^{-10}
RELA	rs3741378	11	65165513	T	1.70×10^{-6}	4.42×10^{-5}	−0.546	0.087	3.41×10^{-10}
HOXC@	rs7297416	12	52729357	C	2.26×10^{-6}	0.011	−0.334	0.065	2.32×10^{-7}

important in cellular metabolism and has been associated with Wolff-Parkinson-White syndrome (MIM 194200), urate levels and chronic kidney disease,^{81,82} and accumulation of cardiac glycogen and left ventricular hypertrophy resembling hypertrophic cardiomyopathy.^{83,84} Chronic kidney disease and left ventricular hypertrophy represent target organ damage related to hypertension, and thus, our findings may provide a link between genetic loci affecting BP and the occurrence of clinically important sequelae. Other traits associated with *PRKAG2* include hemoglobin⁸⁵ and hematocrits.⁸⁶ The *FBN1* locus encodes fibrillin-1 (*FBN1*), a component of elastic fibers in connective tissue, and this gene has been associated with Marfan syndrome (MIM 154700), with systolic and pulse pressure, with aortic stiffness in patients with coronary artery disease (CAD),^{87,88} and with thoracic aortic aneurysms and thoracic aortic dissection.⁸⁹ *CDK6* variants are associated with height (MIM 606255) in a number of studies^{90–92} and are implicated in white blood cell counts in Japanese populations^{93,94} and in African Americans.⁹⁵

Associated Loci with Multiple Genes

The remaining loci (*VCL*, *H19*, *NUCB2*, *SIPA1*, *HOXC4*, and *NFAT5*) contain several genes that we have prioritized

on the basis of variant functionality and the biological rationale of the genes in the BP phenotypes.

We consider vinculin (*VCL*) the strongest biological candidate. Vinculin is a cytoskeletal protein, localized to intercalated discs, and by anchoring thin filaments it is implicated in cardiac force generation. Targeted disruption of vinculin in mice has shown loss of cardiac contractility in embryonic development.⁹⁶ The two other candidates at the *VCL* locus (*AP3M1* and *ADK*) show moderate evidence of functional variation.

The *H19* locus contains three genes. *H19* (*H19*, imprinted maternally expressed transcript [nonprotein coding]) expresses a noncoding RNA. Methylation defects in this gene have been associated with pre-eclampsia in a study with 188 pregnancies⁹⁷ and imprinting syndromes such as Beckwith-Wiedemann syndrome (MIM 130650) and growth retardation disorder Silver-Russell syndrome (MIM 180860).^{98,99} The locus also contains the mitochondrial ribosomal protein L23 (*MRPL23*) and an antisense transcript *MRPL23-AS1*. There are no data suggesting a putative role in BP for the latter, but eQTL analysis identified a putative isoform of *MRPL23* (AK126915).

The *NUCB2* locus contains four genes: *PIK3C2A*, *NUCB2*, *NCR3LG1*, and *KCNJ11*. The product of *NUCB2*,

nucleobindin 2, induces hypertension when intracerebrally administered.¹⁰⁰ It is also involved in the maintenance of calcium blood levels, feeding behavior, water intake, glucose homeostasis, and the release of tumor necrosis factor from vascular endothelial cells by interacting with *ERAP1*.^{101,102} It has been also associated with height in GWASs.^{90–92} The potassium inwardly rectifying channel, encoded by *KCNJ11*, is also a good biological candidate, which has been reported to be associated with hypertension¹⁰³ as well as type 2 diabetes, although the previously reported variants are not in LD with the association reported in this study.

The *SIPA1* locus contains seven genes: *MAP3K11*, *PCNXL3*, *SIPA1*, *RELA*, *KAT5*, *RNASEH2C*, and *AP5B1*. *SIPA1* (signal-induced proliferation-associated 1) is the strongest functional candidate, and the associated SNP encodes a nonsynonymous p.Ser182Phe polymorphism, which is predicted to be deleterious by several bioinformatics tools.³¹ The strongest biological candidate is the adjacent *RELA* (a.k.a. NF- κ B), which forms part of the NF- κ B protein complex and has been shown to modulate angiotensin II-induced hypertension in the paraventricular nucleus.¹⁰⁴

The *HOXC4* locus contains three homeobox genes: *HOXC4*, *HOXC5*, and *HOXC6*. All three genes are closely related and encode transcription factors involved in morphogenesis. A recent trans-ethnic GWAS on blood pressure found a signal related to *HOXC4*,¹⁰⁵ suggesting that homeoboxes may have an association with blood pressure in more ethnicities.

The *NFAT5* locus also contains several candidates, including *NFAT5* (Nuclear factor of activated T cells 5). *NFAT5* is a transcription factor, recently shown to regulate vascular smooth muscle cell (VSMC) modulation.¹⁰⁶ In vivo studies in *NFAT5*^{+/-}*ApoE*^{-/-} mice indicated that *NFAT5* is directly involved in atherosclerotic lesion formation and identified *NFAT5* as a positive regulator of atherosclerotic lesion formation.¹⁰⁷ A variant at this locus has also been associated with serum urate⁸¹ and age at menarche.¹⁰⁸ Small-molecule activators of the adjacent *NQO1* (NAD(P)H:quinone oxidoreductase) have been shown to ameliorate spontaneous hypertension in animal models via modulation of eNOS activity.¹⁰⁹ The Ubiquitin-protein ligase *WWP2* is also a candidate, because it is known to bind and downregulate the epithelial Na(+) channel (ENaC). Mutations in *WWP2* have been shown to result in hypertension.¹¹⁰

Therapeutic Opportunities

Current antihypertensive medications do not show complete efficacy in all subjects and often require combination therapy of three or more drugs to reach a target blood pressure, and ~10% of individuals show limited reduction in blood pressure on all therapeutic regimens.¹¹¹ This highlights the need for additional antihypertensive medications, both to improve efficacy of treatment and reduce the burden of side effects experi-

enced, particularly where combination therapies are concerned. Among the associations reported in this manuscript, there is evidence of association for 15 genes that are either current drug targets or are potentially druggable based on the predicted potential of a protein to be modified by a small-molecule drug in our analyses. Two genes are targeted by currently widely applied drugs. The *KCNJ11* product is targeted by several agents including the antihypertensive drug verapamil (DrugBank ID DB00661) and the glucose-lowering agent glyburide (DrugBank DB01016). *NQO1* is targeted by several marketed anticoagulant drugs, including dicumarol and menadione (DrugBank DB00266 and DB00170). The protein encoded by *RELA* is part of the NF- κ B protein complex, which is considered to be inhibited by several known drugs, including the antihypertensive olmesartan (DrugBank DB00275)¹¹² and the alcohol deterrent drug disulfiram (DrugBank DB00822). A known side effect of disulfiram in presence of alcohol exposure is hypotension, which gives support to a potential role of *RELA* in hypertension.

Nine genes with evidence of association have also previously described small-molecule modulators (mainly inhibitory or binding) based on a query of the ChEMBL database. The genes are listed with the number of tool compounds in parentheses: *PDE1A* (42), *CDK6* (405), *ADK* (483), *MAP3K11* (192), *RELA* (289), *KAT5* (25), and *PIK3C2A* (10). The number of molecules identified may indicate the extent of drug discovery research that has been focused on each target. Many are likely to be the focus of pre-existing drug development in industry that may still be ongoing or terminated.¹¹³ Once molecular properties of these compounds are considered to have a favorable profile, they could be investigated in animal models of hypertension; a large number of compounds have already been characterized: the ChEMBL database reports that 405 molecules show activity against *CDK6*. Most of these *CDK6* inhibitors originate from the published GSK (Glaxosmithkline) kinase inhibitor set.¹¹³ A review of the molecular properties of these molecules in ChEMBL shows that they have similar drug-like features and are likely to be orally bioavailable (based on Lipinski's rule of five compliance¹¹⁴). As we described earlier, association is limited to *CDK6* only, and because many of these drug-like molecules would be suitable for immediate evaluation in animal models of hypertension, this might be a worthwhile experiment, even in the absence of other strong evidence to support the role of *CDK6* in BP.

The 11 loci identified in the present study increase our knowledge on BP-related processes and shed light on plausible candidates at each locus and networks, via eQTL and pathway analysis. It is particularly interesting that most of the genes within the associations are suitable candidates for existing drugs or preclinical compounds. These new observations will help to improve our knowledge on BP and related mechanisms.

Supplemental Data

Supplemental Data include Acknowledgments and six tables and can be found with this article online at <http://www.cell.com/AJHG/>.

Received: September 30, 2013

Accepted: December 20, 2013

Published: February 20, 2014

Web Resources

The URLs for data presented herein are as follows:

ChEMBL, <http://www.ebi.ac.uk/chembl>

DrugBank, <http://www.drugbank.ca>

Online Mendelian Inheritance in Man (OMIM), <http://www.omim.org/>

SNP Annotation and Proxy Search (SNAP), <http://www.broadinstitute.org/mpg/snap/>

UCSC Genome Browser, <http://genome.ucsc.edu>

References

1. Wilson, P.W. (1994). Established risk factors and coronary artery disease: the Framingham Study. *Am. J. Hypertens.* 7, 7S–12S.
2. Lawes, C.M.M., Vander Hoorn, S., and Rodgers, A.; International Society of Hypertension (2008). Global burden of blood-pressure-related disease, 2001. *Lancet* 371, 1513–1518.
3. Levy, D., DeStefano, A.L., Larson, M.G., O'Donnell, C.J., Lifton, R.P., Gavvas, H., Cupples, L.A., and Myers, R.H. (2000). Evidence for a gene influencing blood pressure on chromosome 17. Genome scan linkage results for longitudinal blood pressure phenotypes in subjects from the framingham heart study. *Hypertension* 36, 477–483.
4. Kupper, N., Willemsen, G., Riese, H., Posthuma, D., Boomsma, D.I., and de Geus, E.J. (2005). Heritability of daytime ambulatory blood pressure in an extended twin design. *Hypertension* 45, 80–85.
5. Levy, D., Ehret, G.B., Rice, K., Verwoert, G.C., Launer, L.J., Dehghan, A., Glazer, N.L., Morrison, A.C., Johnson, A.D., Aspelund, T., et al. (2009). Genome-wide association study of blood pressure and hypertension. *Nat. Genet.* 41, 677–687.
6. Padmanabhan, S., Newton-Cheh, C., and Dominiczak, A.F. (2012). Genetic basis of blood pressure and hypertension. *Trends Genet.* 28, 397–408.
7. Franceschini, N., Reiner, A.P., and Heiss, G. (2011). Recent findings in the genetics of blood pressure and hypertension traits. *Am. J. Hypertens.* 24, 392–400.
8. Ganesh, S.K., Tragante, V., Guo, W., Guo, Y., Lanktree, M.B., Smith, E.N., Johnson, T., Castillo, B.A., Barnard, J., Baumert, J., et al.; CARDIOGRAM, METASTROKE; LifeLines Cohort Study (2013). Loci influencing blood pressure identified using a cardiovascular gene-centric array. *Hum. Mol. Genet.* 22, 1663–1678.
9. Ehret, G.B., Munroe, P.B., Rice, K.M., Bochud, M., Johnson, A.D., Chasman, D.I., Smith, A.V., Tobin, M.D., Verwoert, G.C., Hwang, S.J., et al.; International Consortium for Blood Pressure Genome-Wide Association Studies; CARDIOGRAM consortium; CKDGen Consortium; KidneyGen Consortium; EchoGen consortium; CHARGE-HF consortium (2011). Genetic variants in novel pathways influence blood pressure and cardiovascular disease risk. *Nature* 478, 103–109.
10. Newton-Cheh, C., Johnson, T., Gateva, V., Tobin, M.D., Bochud, M., Coin, L., Najjar, S.S., Zhao, J.H., Heath, S.C., Eyheramendy, S., et al.; Wellcome Trust Case Control Consortium (2009). Genome-wide association study identifies eight loci associated with blood pressure. *Nat. Genet.* 41, 666–676.
11. Johnson, T., Gaunt, T.R., Newhouse, S.J., Padmanabhan, S., Tomaszewski, M., Kumari, M., Morris, R.W., Tzoulaki, I., O'Brien, E.T., Poulter, N.R., et al.; Cardiogenics Consortium; Global BPgen Consortium (2011). Blood pressure loci identified with a gene-centric array. *Am. J. Hum. Genet.* 89, 688–700.
12. Zhu, X., Young, J.H., Fox, E., Keating, B.J., Franceschini, N., Kang, S., Tayo, B., Adeyemo, A., Sun, Y.V., Li, Y., et al. (2011). Combined admixture mapping and association analysis identifies a novel blood pressure genetic locus on 5p13: contributions from the CArE consortium. *Hum. Mol. Genet.* 20, 2285–2295.
13. Zhan, W., Cruickshanks, K.J., Klein, B.E.K., Klein, R., Huang, G.-H., Pankow, J.S., Gangnon, R.E., and Tweed, T.S. (2010). Generational differences in the prevalence of hearing impairment in older adults. *Am. J. Epidemiol.* 171, 260–266.
14. Keating, B.J., Tischfield, S., Murray, S.S., Bhangale, T., Price, T.S., Glessner, J.T., Galver, L., Barrett, J.C., Grant, S.F., Farlow, D.N., et al. (2008). Concept, design and implementation of a cardiovascular gene-centric 50 k SNP array for large-scale genomic association studies. *PLoS ONE* 3, e3583.
15. Tomaszewski, M., Debiec, R., Braund, P.S., Nelson, C.P., Hardwick, R., Christofidou, P., Denniff, M., Codd, V., Rafelt, S., van der Harst, P., et al. (2010). Genetic architecture of ambulatory blood pressure in the general population: insights from cardiovascular gene-centric array. *Hypertension* 56, 1069–1076.
16. Clarke, R., Peden, J.F., Hopewell, J.C., Kyriakou, T., Goel, A., Heath, S.C., Parish, S., Barlera, S., Franzosi, M.G., Rust, S., et al.; PROCARDIS Consortium (2009). Genetic variants associated with Lp(a) lipoprotein level and coronary disease. *N. Engl. J. Med.* 361, 2518–2528.
17. IBC 50K CAD Consortium (2011). Large-scale gene-centric analysis identifies novel variants for coronary artery disease. *PLoS Genet.* 7, e1002260.
18. Asselbergs, F.W., Guo, Y., van Iperen, E.P., Sivapalaratnam, S., Tragante, V., Lanktree, M.B., Lange, L.A., Almoguera, B., Appelman, Y.E., Barnard, J., et al.; LifeLines Cohort Study (2012). Large-scale gene-centric meta-analysis across 32 studies identifies multiple lipid loci. *Am. J. Hum. Genet.* 91, 823–838.
19. Cappola, T.P., Li, M., He, J., Ky, B., Gilmore, J., Qu, L., Keating, B., Reilly, M., Kim, C.E., Glessner, J., et al. (2010). Common variants in HSPB7 and FRMD4B associated with advanced heart failure. *Circ Cardiovasc Genet* 3, 147–154.
20. Saxena, R., Elbers, C.C., Guo, Y., Peter, I., Gaunt, T.R., Mega, J.L., Lanktree, M.B., Tare, A., Castillo, B.A., Li, Y.R., et al.; Look AHEAD Research Group; DIAGRAM consortium (2012). Large-scale gene-centric meta-analysis across 39 studies identifies type 2 diabetes loci. *Am. J. Hum. Genet.* 90, 410–425.

21. Guo, Y., Lanktree, M.B., Taylor, K.C., Hakonarson, H., Lange, L.A., and Keating, B.J.; IBC 50K SNP array BMI Consortium (2013). Gene-centric meta-analyses of 108 912 individuals confirm known body mass index loci and reveal three novel signals. *Hum. Mol. Genet.* 22, 184–201.
22. Johnson, A.D., Newton-Cheh, C., Chasman, D.I., Ehret, G.B., Johnson, T., Rose, L., Rice, K., Verwoert, G.C., Launer, L.J., Gudnason, V., et al.; Cohorts for Heart and Aging Research in Genomic Epidemiology Consortium; Global BPgen Consortium; Women's Genome Health Study (2011). Association of hypertension drug target genes with blood pressure and hypertension in 86,588 individuals. *Hypertension* 57, 903–910.
23. de Bakker, P.I.W., Ferreira, M.A.R., Jia, X., Neale, B.M., Raychaudhuri, S., and Voight, B.F. (2008). Practical aspects of imputation-driven meta-analysis of genome-wide association studies. *Hum. Mol. Genet.* 17 (R2), R122–R128.
24. Heap, G.A., Trynka, G., Jansen, R.C., Bruinenberg, M., Swertz, M.A., Dinesen, L.C., Hunt, K.A., Wijmenga, C., Vanheer, D.A., and Franke, L. (2009). Complex nature of SNP genotype effects on gene expression in primary human leucocytes. *BMC Med. Genomics* 2, 1.
25. Lo, K.S., Wilson, J.G., Lange, L.A., Folsom, A.R., Galarneau, G., Ganesh, S.K., Grant, S.F., Keating, B.J., McCarroll, S.A., Mohler, E.R., 3rd., et al. (2011). Genetic association analysis highlights new loci that modulate hematological trait variation in Caucasians and African Americans. *Hum. Genet.* 129, 307–317.
26. Johnson, A.D., Handsaker, R.E., Pulit, S.L., Nizzari, M.M., O'Donnell, C.J., and de Bakker, P.I. (2008). SNAP: a web-based tool for identification and annotation of proxy SNPs using HapMap. *Bioinformatics* 24, 2938–2939.
27. Ridker, P.M., Chasman, D.I., Zee, R.Y., Parker, A., Rose, L., Cook, N.R., and Buring, J.E.; Women's Genome Health Study Working Group (2008). Rationale, design, and methodology of the Women's Genome Health Study: a genome-wide association study of more than 25,000 initially healthy american women. *Clin. Chem.* 54, 249–255.
28. Diercks, G.F., van Boven, A.J., Hillege, H.L., Janssen, W.M., Kors, J.A., de Jong, P.E., Grobbee, D.E., Crijns, H.J., and van Gilst, W.H. (2000). Microalbuminuria is independently associated with ischaemic electrocardiographic abnormalities in a large non-diabetic population. The PREVEND (Prevention of Renal and Vascular ENDstage Disease) study. *Eur. Heart J.* 21, 1922–1927.
29. Stolk, R.P., Rosmalen, J.G., Postma, D.S., de Boer, R.A., Navis, G., Slaets, J.P., Ormel, J., and Wolffenbuttel, B.H. (2008). Universal risk factors for multifactorial diseases: LifeLines: a three-generation population-based study. *Eur. J. Epidemiol.* 23, 67–74.
30. Ward, L.D., and Kellis, M. (2012). HaploReg: a resource for exploring chromatin states, conservation, and regulatory motif alterations within sets of genetically linked variants. *Nucleic Acids Res.* 40 (D1), D930–D934.
31. Wang, K., Li, M., and Hakonarson, H. (2010). ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. *Nucleic Acids Res.* 38, e164.
32. Boyle, A.P., Hong, E.L., Hariharan, M., Cheng, Y., Schaub, M.A., Kasowski, M., Karczewski, K.J., Park, J., Hitz, B.C., Weng, S., et al. (2012). Annotation of functional variation in personal genomes using RegulomeDB. *Genome Res.* 22, 1790–1797.
33. Bernstein, B.E., Birney, E., Dunham, I., Green, E.D., Gunter, C., and Snyder, M.; ENCODE Project Consortium (2012). An integrated encyclopedia of DNA elements in the human genome. *Nature* 489, 57–74.
34. Liu, Y.I., Wise, P.H., and Butte, A.J. (2009). The “etiome”: identification and clustering of human disease etiological factors. *BMC Bioinformatics* 10 (Suppl 2), S14.
35. Talmud, P.J., Drenos, F., Shah, S., Shah, T., Palmen, J., Verzilli, C., Gaunt, T.R., Pallas, J., Lovering, R., Li, K., et al.; ASCOT investigators; NORDIL investigators; BRIGHT Consortium (2009). Gene-centric association signals for lipids and apolipoproteins identified via the HumanCVD BeadChip. *Am. J. Hum. Genet.* 85, 628–642.
36. Ekins, S., Mestres, J., and Testa, B. (2007). In silico pharmacology for drug discovery: applications to targets and beyond. *Br. J. Pharmacol.* 152, 21–37.
37. Volkamer, A., Kuhn, D., Grombacher, T., Rippmann, F., and Rarey, M. (2012). Combining global and local measures for structure-based druggability predictions. *J. Chem. Inf. Model.* 52, 360–372.
38. Göring, H.H., Curran, J.E., Johnson, M.P., Dyer, T.D., Charlesworth, J., Cole, S.A., Jowett, J.B., Abraham, L.J., Rainwater, D.L., Comuzzie, A.G., et al. (2007). Discovery of expression QTLs using large-scale transcriptional profiling in human lymphocytes. *Nat. Genet.* 39, 1208–1216.
39. Idaghdour, Y., Czika, W., Shianna, K.V., Lee, S.H., Visscher, P.M., Martin, H.C., Miclaus, K., Jadallah, S.J., Goldstein, D.B., Wolfinger, R.D., and Gibson, G. (2010). Geographical genomics of human leukocyte gene expression variation in southern Morocco. *Nat. Genet.* 42, 62–67.
40. Emilsson, V., Thorleifsson, G., Zhang, B., Leonardson, A.S., Zink, F., Zhu, J., Carlson, S., Helgason, A., Walters, G.B., Gunnarsdottir, S., et al. (2008). Genetics of gene expression and its effect on disease. *Nature* 452, 423–428.
41. Fehrmann, R.S., Jansen, R.C., Veldink, J.H., Westra, H.J., Arends, D., Bonder, M.J., Fu, J., Deelen, P., Groen, H.J., Smolonska, A., et al. (2011). Trans-eQTLs reveal that independent genetic variants associated with a complex phenotype converge on intermediate genes, with a major role for the HLA. *PLoS Genet.* 7, e1002197.
42. Mehta, D., Heim, K., Herder, C., Carstensen, M., Eckstein, G., Schurmann, C., Homuth, G., Nauck, M., Völker, U., Roden, M., et al. (2013). Impact of common regulatory single-nucleotide variants on gene expression profiles in whole blood. *Eur. J. Hum. Genet.* 21, 48–54.
43. Sasayama, D., Hori, H., Nakamura, S., Miyata, R., Teraishi, T., Hattori, K., Ota, M., Yamamoto, N., Higuchi, T., Amano, N., and Kunugi, H. (2013). Identification of single nucleotide polymorphisms regulating peripheral blood mRNA expression with genome-wide significance: an eQTL study in the Japanese population. *PLoS ONE* 8, e54967.
44. Dixon, A.L., Liang, L., Moffatt, M.F., Chen, W., Heath, S., Wong, K.C., Taylor, J., Burnett, E., Gut, I., Farrall, M., et al. (2007). A genome-wide association study of global gene expression. *Nat. Genet.* 39, 1202–1207.
45. Liang, L., Morar, N., Dixon, A.L., Lathrop, G.M., Abecasis, G.R., Moffatt, M.F., and Cookson, W.O. (2013). A cross-platform analysis of 14,177 expression quantitative trait loci derived from lymphoblastoid cell lines. *Genome Res.* 23, 716–726.
46. Stranger, B.E., Nica, A.C., Forrest, M.S., Dimas, A., Bird, C.P., Beazley, C., Ingle, C.E., Dunning, M., Flicek, P., Koller, D.,

- et al. (2007). Population genomics of human gene expression. *Nat. Genet.* 39, 1217–1224.
47. Kwan, T., Benovoy, D., Dias, C., Gurd, S., Provencher, C., Beaulieu, P., Hudson, T.J., Sladek, R., and Majewski, J. (2008). Genome-wide analysis of transcript isoform variation in humans. *Nat. Genet.* 40, 225–231.
48. Dimas, A.S., Deutsch, S., Stranger, B.E., Montgomery, S.B., Borel, C., Attar-Cohen, H., Ingle, C., Beazley, C., Gutierrez Arcelus, M., Sekowska, M., et al. (2009). Common regulatory variation impacts gene expression in a cell type-dependent manner. *Science* 325, 1246–1250.
49. Cusanovich, D.A., Billstrand, C., Zhou, X., Chavarria, C., De Leon, S., Michelini, K., Pai, A.A., Ober, C., and Gilad, Y. (2012). The combination of a genome-wide association study of lymphocyte count and analysis of gene expression data reveals novel asthma candidate genes. *Hum. Mol. Genet.* 21, 2111–2123.
50. Grundberg, E., Small, K.S., Hedman, A.K., Nica, A.C., Buil, A., Keildson, S., Bell, J.T., Yang, T.P., Meduri, E., Barrett, A., et al.; Multiple Tissue Human Expression Resource (MuTHER) Consortium (2012). Mapping cis- and trans-regulatory effects across multiple tissues in twins. *Nat. Genet.* 44, 1084–1089.
51. Mangravite, L.M., Engelhardt, B.E., Medina, M.W., Smith, J.D., Brown, C.D., Chasman, D.I., Mecham, B.H., Howie, B., Shim, H., Naidoo, D., et al. (2013). A statin-dependent QTL for GATM expression is associated with statin-induced myopathy. *Nature* 502, 377–380.
52. Fairfax, B.P., Makino, S., Radhakrishnan, J., Plant, K., Leslie, S., Dillthey, A., Ellis, P., Langford, C., Vannberg, F.O., and Knight, J.C. (2012). Genetics of gene expression in primary immune cells identifies cell type-specific master regulators and roles of HLA alleles. *Nat. Genet.* 44, 502–510.
53. Murphy, A., Chu, J.H., Xu, M., Carey, V.J., Lazarus, R., Liu, A., Szefer, S.J., Strunk, R., Demuth, K., Castro, M., et al. (2010). Mapping of numerous disease-associated expression polymorphisms in primary peripheral blood CD4+ lymphocytes. *Hum. Mol. Genet.* 19, 4745–4757.
54. Heinzen, E.L., Ge, D., Cronin, K.D., Maia, J.M., Shianna, K.V., Gabriel, W.N., Welsh-Bohmer, K.A., Hulette, C.M., Denny, T.N., and Goldstein, D.B. (2008). Tissue-specific genetic control of splicing: implications for the study of complex traits. *PLoS Biol.* 6, e1.
55. Zeller, T., Wild, P., Szymczak, S., Rotival, M., Schillert, A., Castagne, R., Maouche, S., Germain, M., Lackner, K., Rossman, H., et al. (2010). Genetics and beyond—the transcriptome of human monocytes and disease susceptibility. *PLoS ONE* 5, e10693.
56. Barreiro, L.B., Tailleux, L., Pai, A.A., Gicquel, B., Marionni, J.C., and Gilad, Y. (2012). Deciphering the genetic architecture of variation in the immune response to *Mycobacterium tuberculosis* infection. *Proc. Natl. Acad. Sci. USA* 109, 1204–1209.
57. Greenawalt, D.M., Dobrin, R., Chudin, E., Hatoum, I.J., Suver, C., Beaulaurier, J., Zhang, B., Castro, V., Zhu, J., Sieberts, S.K., et al. (2011). A survey of the genetics of stomach, liver, and adipose gene expression from a morbidly obese cohort. *Genome Res.* 21, 1008–1016.
58. Kompass, K.S., and Witte, J.S. (2011). Co-regulatory expression quantitative trait loci mapping: method and application to endometrial cancer. *BMC Med. Genomics* 4, 6.
59. Li, Q., Seo, J.H., Stranger, B., McKenna, A., Pe'er, I., Laframboise, T., Brown, M., Tyekucheva, S., and Freedman, M.L. (2013). Integrative eQTL-based analyses reveal the biology of breast cancer risk loci. *Cell* 152, 633–641.
60. Webster, J.A., Gibbs, J.R., Clarke, J., Ray, M., Zhang, W., Holmans, P., Rohrer, K., Zhao, A., Marlowe, L., Kaleem, M., et al.; NACC-Neuropathology Group (2009). Genetic control of human brain transcript expression in Alzheimer disease. *Am. J. Hum. Genet.* 84, 445–458.
61. Zou, F., Chai, H.S., Younkin, C.S., Allen, M., Crook, J., Pankratz, V.S., Carrasquillo, M.M., Rowley, C.N., Nair, A.A., Middha, S., et al.; Alzheimer's Disease Genetics Consortium (2012). Brain expression genome-wide association study (eGWAS) identifies human disease-associated variants. *PLoS Genet.* 8, e1002707.
62. Colantuoni, C., Lipska, B.K., Ye, T., Hyde, T.M., Tao, R., Leek, J.T., Colantuoni, E.A., Elkahoul, A.G., Herman, M.M., Weinberger, D.R., and Kleinman, J.E. (2011). Temporal dynamics and genetic control of transcription in the human prefrontal cortex. *Nature* 478, 519–523.
63. Liu, C., Cheng, L., Badner, J.A., Zhang, D., Craig, D.W., Redman, M., and Gershon, E.S. (2010). Whole-genome association mapping of gene expression in the human prefrontal cortex. *Mol. Psychiatry* 15, 779–784.
64. Gibbs, J.R., van der Brug, M.P., Hernandez, D.G., Traynor, B.J., Nalls, M.A., Lai, S.L., Arepalli, S., Dillman, A., Rafferty, I.P., Troncoso, J., et al. (2010). Abundant quantitative trait loci exist for DNA methylation and gene expression in human brain. *PLoS Genet.* 6, e1000952.
65. Zhang, B., Gaiteri, C., Bodea, L.-G., Wang, Z., McElwee, J., Podtelezhnikov, A.A., Zhang, C., Xie, T., Tran, L., Dobrin, R., et al. (2013). Integrated systems approach identifies genetic nodes and networks in late-onset Alzheimer's disease. *Cell* 153, 707–720.
66. Schadt, E.E., Molony, C., Chudin, E., Hao, K., Yang, X., Lum, P.Y., Kasarskis, A., Zhang, B., Wang, S., Suver, C., et al. (2008). Mapping the genetic architecture of gene expression in human liver. *PLoS Biol.* 6, e107.
67. Innocenti, F., Cooper, G.M., Stanaway, I.B., Gamazon, E.R., Smith, J.D., Mirkov, S., Ramirez, J., Liu, W., Lin, Y.S., Moloney, C., et al. (2011). Identification, replication, and functional fine-mapping of expression quantitative trait loci in primary human liver tissue. *PLoS Genet.* 7, e1002078.
68. Grundberg, E., Kwan, T., Ge, B., Lam, K.C., Koka, V., Kindmark, A., Mallmin, H., Dias, J., Verlaan, D.J., Ouimet, M., et al. (2009). Population genomics in a disease targeted primary cell model. *Genome Res.* 19, 1942–1952.
69. Kabachiev, B., and Silverberg, M.S. (2013). Expression quantitative trait loci analysis identifies associations between genotype and gene expression in human intestine. *Gastroenterology* 144, 1488–1496, e1–e3.
70. Hao, K., Bossé, Y., Nickle, D.C., Paré, P.D., Postma, D.S., Laviolette, M., Sandford, A., Hackett, T.L., Daley, D., Hogg, J.C., et al. (2012). Lung eQTLs to help reveal the molecular underpinnings of asthma. *PLoS Genet.* 8, e1003029.
71. Ding, J., Gudjonsson, J.E., Liang, L., Stuart, P.E., Li, Y., Chen, W., Weichenthal, M., Ellinghaus, E., Franke, A., Cookson, W., et al. (2010). Gene expression in skin and lymphoblastoid cells: Refined statistical method reveals extensive overlap in cis-eQTL signals. *Am. J. Hum. Genet.* 87, 779–789.
72. Huang, R.S., Gamazon, E.R., Ziliak, D., Wen, Y., Im, H.K., Zhang, W., Wing, C., Duan, S., Bleibel, W.K., Cox, N.J., and Dolan, M.E. (2011). Population differences in microRNA

- expression and biological implications. *RNA Biol.* 8, 692–701.
73. Rantalainen, M., Herrera, B.M., Nicholson, G., Bowden, R., Wills, Q.F., Min, J.L., Neville, M.J., Barrett, A., Allen, M., Rayner, N.W., et al. (2011). MicroRNA expression in abdominal and gluteal adipose tissue is associated with mRNA expression levels and partly genetically driven. *PLoS ONE* 6, e27338.
 74. Newton-Cheh, C., Larson, M.G., Vasan, R.S., Levy, D., Bloch, K.D., Surti, A., Guiducci, C., Kathiresan, S., Benjamin, E.J., Struck, J., et al. (2009). Association of common variants in NPPA and NPPB with circulating natriuretic peptides and blood pressure. *Nat. Genet.* 41, 348–353.
 75. Salvi, E., Kutalik, Z., Glorioso, N., Benaglio, P., Frau, F., Kuznetsova, T., Arima, H., Hoggart, C., Tichet, J., Nikitin, Y.P., et al. (2012). Genomewide association study using a high-density single nucleotide polymorphism array and case-control design identifies a novel essential hypertension susceptibility locus in the promoter region of endothelial NO synthase. *Hypertension* 59, 248–255.
 76. Wain, L.V., Verwoert, G.C., O'Reilly, P.F., Shi, G., Johnson, T., Johnson, A.D., Bochud, M., Rice, K.M., Henneman, P., Smith, A.V., et al.; LifeLines Cohort Study; EchoGen consortium; AortaGen Consortium; CHARGE Consortium Heart Failure Working Group; KidneyGen consortium; CKDGen consortium; Cardiogenics consortium; CardioGram (2011). Genome-wide association study identifies six new loci influencing pulse pressure and mean arterial pressure. *Nat. Genet.* 43, 1005–1011.
 77. Padmanabhan, S., Melander, O., Johnson, T., Di Blasio, A.M., Lee, W.K., Gentilini, D., Hastie, C.E., Menni, C., Monti, M.C., Delles, C., et al.; Global BPgen Consortium (2010). Genome-wide association study of blood pressure extremes identifies variant near UMOD associated with hypertension. *PLoS Genet.* 6, e1001177.
 78. Giachini, F.R., Lima, V.V., Carneiro, F.S., Tostes, R.C., and Webb, R.C. (2011). Decreased cGMP level contributes to increased contraction in arteries from hypertensive rats: role of phosphodiesterase 1. *Hypertension* 57, 655–663.
 79. Schermlay, R.T., Pullamsetti, S.S., Kwapiszewska, G., Dumitrascu, R., Tian, X., Weissmann, N., Ghofrani, H.A., Kaulen, C., Dunkern, T., Schudt, C., et al. (2007). Phosphodiesterase 1 upregulation in pulmonary arterial hypertension: target for reverse-remodeling therapy. *Circulation* 115, 2331–2339.
 80. Zhu, F., Sun, Y., Wang, M., Ma, S., Chen, X., Cao, A., Chen, F., Qiu, Y., and Liao, Y. (2011). Correlation between HLA-DRB1, HLA-DQB1 polymorphism and autoantibodies against angiotensin AT(1) receptors in Chinese patients with essential hypertension. *Clin. Cardiol.* 34, 302–308.
 81. Köttgen, A., Albrecht, E., Teumer, A., Vitart, V., Krumsiek, J., Hundertmark, C., Pistis, G., Ruggiero, D., O'Seaghdha, C.M., Haller, T., et al.; LifeLines Cohort Study; CARDIoGRAM Consortium; DIAGRAM Consortium; ICBP Consortium; MAGIC Consortium (2013). Genome-wide association analyses identify 18 new loci associated with serum urate concentrations. *Nat. Genet.* 45, 145–154.
 82. Köttgen, A., Pattaro, C., Böger, C.A., Fuchsberger, C., Olden, M., Glazer, N.L., Parsa, A., Gao, X., Yang, Q., Smith, A.V., et al. (2010). New loci associated with kidney function and chronic kidney disease. *Nat. Genet.* 42, 376–384.
 83. Gollob, M.H., Green, M.S., Tang, A.S., Gollob, T., Karibe, A., Ali Hassan, A.S., Ahmad, F., Lozado, R., Shah, G., Fanana-pazir, L., et al. (2001). Identification of a gene responsible for familial Wolff-Parkinson-White syndrome. *N. Engl. J. Med.* 344, 1823–1831.
 84. Arad, M., Maron, B.J., Gorham, J.M., Johnson, W.H., Jr., Saul, J.P., Perez-Atayde, A.R., Spirito, P., Wright, G.B., Kanter, R.J., Seidman, C.E., and Seidman, J.G. (2005). Glycogen storage diseases presenting as hypertrophic cardiomyopathy. *N. Engl. J. Med.* 352, 362–372.
 85. van der Harst, P., Zhang, W., Mateo Leach, I., Rendon, A., Verweij, N., Sehmi, J., Paul, D.S., Elling, U., Allayee, H., Li, X., et al. (2012). Seventy-five genetic loci influencing the human red blood cell. *Nature* 492, 369–375.
 86. Ganesh, S.K., Zakai, N.A., van Rooij, F.J., Soranzo, N., Smith, A.V., Nalls, M.A., Chen, M.H., Kottgen, A., Glazer, N.L., Dehghan, A., et al. (2009). Multiple loci influence erythrocyte phenotypes in the CHARGE Consortium. *Nat. Genet.* 41, 1191–1198.
 87. Powell, J.T., Turner, R.J., Henney, A.M., Miller, G.J., and Humphries, S.E. (1997). An association between arterial pulse pressure and variation in the fibrillin-1 gene. *Heart* 78, 396–398.
 88. Medley, T.L., Cole, T.J., Gatzka, C.D., Wang, W.Y.S., Dart, A.M., and Kingwell, B.A. (2002). Fibrillin-1 genotype is associated with aortic stiffness and disease severity in patients with coronary artery disease. *Circulation* 105, 810–815.
 89. Lemaire, S.A., McDonald, M.-L.N., Guo, D.C., Russell, L., Miller, C.C., 3rd, Johnson, R.J., Bekheirnia, M.R., Franco, L.M., Nguyen, M., Pyeritz, R.E., et al. (2011). Genome-wide association study identifies a susceptibility locus for thoracic aortic aneurysms and aortic dissections spanning FBN1 at 15q21.1. *Nat. Genet.* 43, 996–1000.
 90. Lettre, G., Jackson, A.U., Gieger, C., Schumacher, F.R., Berndt, S.I., Sanna, S., Eyheramendy, S., Voight, B.F., Butler, J.L., Guiducci, C., et al.; Diabetes Genetics Initiative; FUSION; KORA; Prostate, Lung Colorectal and Ovarian Cancer Screening Trial; Nurses' Health Study; SardiNIA (2008). Identification of ten loci associated with height highlights new biological pathways in human growth. *Nat. Genet.* 40, 584–591.
 91. Lango Allen, H., Estrada, K., Lettre, G., Berndt, S.I., Weedon, M.N., Rivadeneira, F., Willer, C.J., Jackson, A.U., Vedantam, S., Raychaudhuri, S., et al. (2010). Hundreds of variants clustered in genomic loci and biological pathways affect human height. *Nature* 467, 832–838.
 92. Berndt, S.I., Gustafsson, S., Mägi, R., Ganna, A., Wheeler, E., Feitosa, M.F., Justice, A.E., Monda, K.L., Croteau-Chonka, D.C., Day, F.R., et al. (2013). Genome-wide meta-analysis identifies 11 new loci for anthropometric traits and provides insights into genetic architecture. *Nat. Genet.* 45, 501–512.
 93. Kamatani, Y., Matsuda, K., Okada, Y., Kubo, M., Hosono, N., Daigo, Y., Nakamura, Y., and Kamatani, N. (2010). Genome-wide association study of hematological and biochemical traits in a Japanese population. *Nat. Genet.* 42, 210–215.
 94. Okada, Y., Hirota, T., Kamatani, Y., Takahashi, A., Ohmiya, H., Kumasaka, N., Higasa, K., Yamaguchi-Kabata, Y., Hosono, N., Nalls, M.A., et al. (2011). Identification of nine novel loci associated with white blood cell subtypes in a Japanese population. *PLoS Genet.* 7, e1002067.
 95. Reiner, A.P., Lettre, G., Nalls, M.A., Ganesh, S.K., Mathias, R., Austin, M.A., Dean, E., Arepalli, S., Britton, A., Chen, Z., et al. (2011). Genome-wide association study of white blood cell count in 16,388 African Americans: the continental origins

- and genetic epidemiology network (COGENT). *PLoS Genet.* 7, e1002108.
96. Xu, W., Baribault, H., and Adamson, E.D. (1998). Vinculin knockout results in heart and brain defects during embryonic development. *Development* 125, 327–337.
 97. Yu, L., Chen, M., Zhao, D., Yi, P., Lu, L., Han, J., Zheng, X., Zhou, Y., and Li, L. (2009). The H19 gene imprinting in normal pregnancy and pre-eclampsia. *Placenta* 30, 443–447.
 98. Sparago, A., Cerrato, F., Vernucci, M., Ferrero, G.B., Silengo, M.C., and Riccio, A. (2004). Microdeletions in the human H19 DMR result in loss of IGF2 imprinting and Beckwith-Wiedemann syndrome. *Nat. Genet.* 36, 958–960.
 99. Blik, J., Terhal, P., van den Bogaard, M.-J., Maas, S., Hamel, B., Salieb-Beugelaar, G., Simon, M., Letteboer, T., van der Smagt, J., Kroes, H., and Mannens, M. (2006). Hypomethylation of the H19 gene causes not only Silver-Russell syndrome (SRS) but also isolated asymmetry or an SRS-like phenotype. *Am. J. Hum. Genet.* 78, 604–614.
 100. Angelone, T., Filice, E., Pasqua, T., Amodio, N., Galluccio, M., Montesanti, G., Quintieri, A.M., and Cerra, M.C. (2013). Nesfatin-1 as a novel cardiac peptide: identification, functional characterization, and protection against ischemia/reperfusion injury. *Cell. Mol. Life Sci.* 70, 495–509.
 101. Islam, A., Adamik, B., Hawari, F.I., Ma, G., Rouhani, F.N., Zhang, J., and Levine, S.J. (2006). Extracellular TNFR1 release requires the calcium-dependent formation of a nucleobindin 2-ARTS-1 complex. *J. Biol. Chem.* 281, 6860–6873.
 102. Oh-I, S., Shimizu, H., Satoh, T., Okada, S., Adachi, S., Inoue, K., Eguchi, H., Yamamoto, M., Imaki, T., Hashimoto, K., et al. (2006). Identification of nesfatin-1 as a satiety molecule in the hypothalamus. *Nature* 443, 709–712.
 103. Kokubo, Y., Tomoike, H., Tanaka, C., Banno, M., Okuda, T., Inamoto, N., Kamide, K., Kawano, Y., and Miyata, T. (2006). Association of sixty-one non-synonymous polymorphisms in forty-one hypertension candidate genes with blood pressure variation and hypertension. *Hypertens. Res.* 29, 611–619.
 104. Cardinale, J.P., Sriramula, S., Mariappan, N., Agarwal, D., and Francis, J. (2012). Angiotensin II-induced hypertension is modulated by nuclear factor- κ B in the paraventricular nucleus. *Hypertension* 59, 113–121.
 105. Franceschini, N., Fox, E., Zhang, Z., Edwards, T.L., Nalls, M.A., Sung, Y.J., Tayo, B.O., Sun, Y.V., Gottesman, O., Adeyemo, A., et al.; Asian Genetic Epidemiology Network Consortium (2013). Genome-wide association analysis of blood-pressure traits in African-ancestry individuals reveals common associated genes in African and non-African populations. *Am. J. Hum. Genet.* 93, 545–554.
 106. Halterman, J.A., Kwon, H.M., Zargham, R., Bortz, P.D., and Wamhoff, B.R. (2011). Nuclear factor of activated T cells 5 regulates vascular smooth muscle cell phenotypic modulation. *Arterioscler. Thromb. Vasc. Biol.* 31, 2287–2296.
 107. Halterman, J.A., Kwon, H.M., Leitinger, N., and Wamhoff, B.R. (2012). NFAT5 expression in bone marrow-derived cells enhances atherosclerosis and drives macrophage migration. *Front. Physiol.* 3, 313.
 108. Elks, C.E., Perry, J.R., Sulem, P., Chasman, D.I., Franceschini, N., He, C., Lunetta, K.L., Visser, J.A., Byrne, E.M., Cousminer, D.L., et al.; GIANT Consortium (2010). Thirty new loci for age at menarche identified by a meta-analysis of genome-wide association studies. *Nat. Genet.* 42, 1077–1085.
 109. Kim, Y.-H., Hwang, J.H., Noh, J.-R., Gang, G.-T., Kim, H., Son, H.Y., Kwak, T.H., Shong, M., Lee, I.K., and Lee, C.H. (2011). Activation of NAD(P)H:quinone oxidoreductase ameliorates spontaneous hypertension in an animal model via modulation of eNOS activity. *Cardiovasc. Res.* 91, 519–527.
 110. McDonald, F.J., Western, A.H., McNeil, J.D., Thomas, B.C., Olson, D.R., and Snyder, P.M. (2002). Ubiquitin-protein ligase WWP2 binds to and downregulates the epithelial Na(+) channel. *Am. J. Physiol. Renal Physiol.* 283, F431–F436.
 111. Persell, S.D. (2011). Prevalence of resistant hypertension in the United States, 2003–2008. *Hypertension* 57, 1076–1080.
 112. Cvek, B., and Dvorak, Z. (2007). Targeting of nuclear factor-kappaB and proteasome by dithiocarbamate complexes with metals. *Curr. Pharm. Des.* 13, 3155–3167.
 113. Dranchak, P., MacArthur, R., Guha, R., Zuercher, W.J., Drewry, D.H., Auld, D.S., and Inglese, J. (2013). Profile of the GSK published protein kinase inhibitor set across ATP-dependent and-independent luciferases: implications for reporter-gene assays. *PLoS ONE* 8, e57888.
 114. Lipinski, C.A., Lombardo, F., Dominy, B.W., and Feeney, P.J. (2001). Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings. *Adv. Drug Deliv. Rev.* 46, 3–26.